

municipality of Escuinapa, Sinaloa (CONABIO 2018. Naturalista. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México, D.F., Mexico; <http://www.naturalista.mx/observations/1832317>; accessed 4 February 2018), and ca. 30 airline km northwest of the closest reported site in Nayarit near Mesa de Pedro y Pablo, Municipality of Acaponeta (National Museum of Natural History, Smithsonian Institution [USNM] 46252; Legler and Vogt 2013. *The Turtles of Mexico: Land and Freshwater Forms*. University of California Press, Berkeley, California. 416 pp.). The turtle was found around 0500 h while it was active on wet forest litter in tropical deciduous forest after heavy rainfall the previous night. We thank the people of Huajicori for their help in the field during the program *Community Monitors to the Natural Patrimony of Sierra Madre Occidental de Nayarit*, which was funded by the project “Payments for Environmental Services” from Consultoría Forestal y Servicios Ambientales Nayarit through the Comisión Nacional Forestal (CONAFOR).

**JESÚS A. LOC BARRAGÁN**, Asociación para la Investigación y Conservación de Anfibios y Reptiles (AICAR) A.C., and Consultoría Forestal y Servicios Ambientales Nayarit S.C. (e-mail: [biolocbarragan@gmail.com](mailto:biolocbarragan@gmail.com)); **HOMERO QUINTERA CASTAÑEDA**, Consultoría Forestal y Servicios Ambientales Nayarit S.C.; **CARLOS DEL VILLAR RODRÍGUEZ**, Alianza Regional para la Conservación de las Aves de Nayarit, A.C. and Consultoría Forestal y Servicios Ambientales Nayarit S.C.; **GUILLERMO A. WOOLRICH-PIÑA**, Laboratorio de Zoología, División de Biología, Subdirección de Investigación y Posgrado, Instituto Tecnológico Superior de Zacapoaxtla, Puebla, Mexico.

**TRACHEMYS SCRIPTA (Pond Slider)**. MEXICO: COAHUILA: MUNICIPALITY OF ZARAGOZA: Zaragoza, Rio Escondido at the bridge along Mexico Hwy 29 (28.48516°N, 100.91707°W; WGS 84), 355 m elev. 10 March 2018. Sean P. Graham, Mark Black, Mark Herr, and Tomas Hernandez. Verified by Toby Hibbitts. James Scud-day Vertebrate Collections, Sul Ross State University (SRSU-D 131; photo voucher). First municipality record located ca. 80 km northeast of the closest reported locality in Coahuila at Múzquiz, Municipality of Múzquiz. This specimen was consistent with the taxon *Trachemys scripta elegans*. Although it is possible this population could be the result of a translocation, wild populations are known from downstream in the Rio Grande drainage. (Lemos-Espinal and Smith 2007. *Anfibios y Reptiles del Estado de Coahuila, México/Amphibians and Reptiles of the State of Coahuila, Mexico*. UNAM, CONABIO, México, D.F. 613 pp.). This research was conducted on behalf of the Maderas del Carmen Flora and Fauna Protection Area under SEMARNAT collecting permit SGPA/DGVS/00047/18 issued to DL, with SPG listed as a sub-permittee. We thank B. Pat, B. McKinney, J. Villalobos, H. Sotelo Gallardo, and other members of El Carmen conservation project for logistical support.

**TOMAS O. HERNANDEZ**, Department of Biology, Midland College, Midland, Texas 79705, USA; **MARK W. HERR**, Division of Herpetology, University of Kansas Natural History Museum, 1345 Jayhawk Blvd., Lawrence, Kansas 66045, USA (e-mail: [mwherr@gmail.com](mailto:mwherr@gmail.com)); **ALEJANDRO ESPINOSA**, CEMEX El Carmen Nature Reserve, Constitucion 444 Pte. Monterrey, Nuevo León, México; **DAVID LAZCANO**, Universidad Autónoma de Nuevo León, Facultad de Ciencias Biológicas, Laboratorio de Herpetología, Apartado Postal 513, San Nicolas de los Garza, Nuevo León, C.P. 66450, México (e-mail: [imantodes52@hotmail.com](mailto:imantodes52@hotmail.com)); **MARK BLACK** and **SEAN P. GRAHAM**, Department of Biology, Geology, and Physical Sciences, Sul Ross State University, Alpine, Texas 79830, USA.

**TRACHEMYS SCRIPTA ELEGANS (Red-eared Slider)**. USA: NEW MEXICO: CURRY Co.: Clovis, Greene Acres Park Lake (34.41850°N, 103.20644°W; WGS 84), 1300 m elev. 17 April 2013. J. N. Stuart. University of Kansas Digital Archives (KUDA 12156; photo voucher). Clovis, Hillcrest City Park, fishing pond (34.41016°N, 103.18544°W; WGS 84), 1456 m elev. 21 April 2015. J. N. Stuart. KUDA 12157 (photo voucher). Ned Houk Memorial Park, ca. 13 km NNE of Clovis (center), fishing pond (34.51606°N, 103.17080°W; WGS 84), 1298 m elev. 21 April 2015. J. N. Stuart. KUDA 12158 (photo voucher). All verified by Leland J. S. Pierce. Multiple individuals seen at each of these three sites. Reported as an introduced species in Curry County by Painter et al. (2017. *West. Wildl.* 4:29–60) but without locality data. The species has been introduced to the few perennial water bodies (all isolated ponds) in the county and is potentially established there. Nearest other confirmed record is ca. 97 km to the north-northwest in the Canadian River near Logan, Quay County, where it is native (Degenhardt and Christiansen. 1974. *Southwest. Nat.* 19:21–46; Degenhardt et al. 1996. *Amphibians and Reptiles of New Mexico*. University of New Mexico Press, Albuquerque, New Mexico. 431 pp.).

**JAMES N. STUART**, New Mexico Department of Game and Fish, P.O. Box 25112, Santa Fe, New Mexico 87504, USA; e-mail: [james.stuart@state.nm.us](mailto:james.stuart@state.nm.us).

## SQUAMATA — LIZARDS

**BRACHYMELES BONITAE (Stub-limbed Burrowing Skink)**. PHILIPPINES: LUZON ISLAND: CAMARINES NORTE PROVINCE: MUNICIPALITY OF LABO: Barangay of Tulay na Lupa, Mt. Labo (14.0802°N, 122.77815°E; WGS 84), 264 m elev. 21–23 May 2017. J. B. Fernandez, E. Schaper, and E. Freitas. Verified by Luke J. Welton. Sam Noble Oklahoma Museum of Natural History (OMNH 45575, 45577, 45578). Collected between 0800 and 1430 h from rotting tree stumps among young herbaceous plants and roots in coconut plantation or beneath piles of decaying coconut husks, leaves, and roots in secondary growth forest. New municipality and province record, and first confirmed record of a member of *B. bonitae* complex from Bicol Peninsula, southeastern Luzon Island (Davis et al. 2014. *J. Herpetol.* 48:480–494; Davis et al. 2016. *Zootaxa* 4132:30–43), extending distribution 113.6 km SE in neighboring province. True *B. bonitae* restricted previously to Central Luzon and Polillo Island (Davis et al. 2014, *op. cit.*). Fieldwork supported by NSF IOS 1353683 to CDS and NSF IOS 1353703 to PJB. Fieldwork was conducted under the Memorandum of Agreement with the BMB of the Philippines (2015–2020) and Gratuitous Permits to Collect No. 247.

**ERIKA G. SCHAPER** (e-mail: [erschaper@clarku.edu](mailto:erschaper@clarku.edu)), **PHILIP J. BERGMANN** (e-mail: [pbergmann@clarku.edu](mailto:pbergmann@clarku.edu)), and **GEN MORINAGA**, Department of Biology, Clark University, 950 Main Street, Worcester, Massachusetts 01610, USA (e-mail: [gmorinaga@clarku.edu](mailto:gmorinaga@clarku.edu)); **JASON FERNANDEZ**, Imus, Cavite Province, Luzon Island, Philippines (e-mail: [jasonfernandez509@gmail.com](mailto:jasonfernandez509@gmail.com)); **ELYSE S. FREITAS** (e-mail: [efreitas@ou.edu](mailto:efreitas@ou.edu)) and **CAMERON D. SILER** (e-mail: [camsiler@ou.edu](mailto:camsiler@ou.edu)), Sam Noble Oklahoma Museum of Natural History, University of Oklahoma, 2401 Chautauqua Ave., Norman, Oklahoma 73072, USA.

**HEMIDACTYLUS TURCICUS (Mediterranean Gecko)**. USA: NEW MEXICO: EDDY Co.: Whites City campground, Carlsbad (34.17527°N, 104.38039°W; WGS 84), ca. 1113 m elev. 20 June 2018. Korry J. Waldon and Thanchira Suriyamongkol. Verified by Carl J. Franklin. Amphibian and Reptile Diversity Research